

HPV (human papillomavirus) is the most common sexually transmitted infection in the United States. Over 100 different types of HPV have been identified.

Possible Signs and Symptoms

- Almost all types of HPV do not produce symptoms.
- If symptoms are present, they may appear within weeks or months after infection

Some strands of HPV cause genital warts. Signs of genital warts may be:

- Soft, pink or flesh-colored bumps or swellings on or near the genitals or anus or upper thigh area
- Cauliflower like growths on or near genitals or anus or upper thigh area

Transmission

- Oral, anal, or vaginal sex
- Infected mother to newborn during vaginal birth

Prevention

- Abstain from oral, anal, and vaginal sex
- Consistent, correct use of latex or polyurethane condoms or barriers during oral, anal, or vaginal sex
- Regular, routine pelvic examinations that include a pap smear test (women)
- Regular, routine health care examinations, including STD tests (men and women)
- If a partner is infected, avoid sexual contact if warts are present and use latex or polyurethane condoms or barriers at other times

For more information about STDs or about testing and treatment in New Jersey, please call the HIV/STD hotline or visit our web site:

1~800~624~2377

http://www.nj.gov/health/cd/stdhome.htm

Treatment

- Examination and tests by a health care provider to diagnose and determine best course of treatment
- There is no cure for HPV, however warts may be removed
- Medications may be warranted in the case of pre-cancerous changes that occur as a result of infection
- Vaccines are currently in production to prevent HPV

Complications/Effects if Left Untreated

- Abnormal pap smear tests
- Increased risk for cervical cancer and/or penile cancer
- Increased risk for HIV infection

If left untreated during pregnancy:

• Can be passed to infant during delivery, may cause warts in throat or voice box of baby

If You Have HPV...

- Refrain from sexual contact (oral, anal, or vaginal sex) if warts are present and use latex or polyurethane condoms or barriers at other times
- Learn about HPV and seek support from local groups or web sites
- Seek medical attention, including STD tests
- Take all medications prescribed and as directed
- Make sure all recent sex partners are notified if you have HPV and talk to new ones about it
- Annual STD screenings, including a pap smear test, if you are, or have ever been, sexually active
- If you are pregnant, get tested for STDs



